Beatriz Castro Dias Cuyabano<sup>1</sup>, Gabriel Rovere<sup>1,2</sup> & Peter Sørensen<sup>1</sup>



Beatriz Castro Dias Cuyabano<sup>1</sup>, Gabriel Rovere<sup>1,2</sup> & Peter Sørensen<sup>1</sup>

<sup>1</sup>Center for Quantitative Genetics and Genomics, Department of Molecular Biology and Genetics, Aarhus University (DK)
<sup>2</sup>Department of Animal Science/Epidemiology & Biostatistics, Michigan State University (USA)

• Genomic models have their likelihood misspecified to that of the true model that generates a phenotype.



Beatriz Castro Dias Cuyabano<sup>1</sup>, Gabriel Rovere<sup>1,2</sup> & Peter Sørensen<sup>1</sup>

- Genomic models have their likelihood misspecified to that of the true model that generates a phenotype.
- Misspecification: SNP-derived covariance structure, instead of QTL-derived.



Beatriz Castro Dias Cuyabano<sup>1</sup>, Gabriel Rovere<sup>1,2</sup> & Peter Sørensen<sup>1</sup>

- Genomic models have their likelihood misspecified to that of the true model that generates a phenotype.
- Misspecification: SNP-derived covariance structure, instead of QTL-derived.
- Representation of covariance structure of the data may be incorrect.



Beatriz Castro Dias Cuyabano<sup>1</sup>, Gabriel Rovere<sup>1,2</sup> & Peter Sørensen<sup>1</sup>

- Genomic models have their likelihood misspecified to that of the true model that generates a phenotype.
- Misspecification: SNP-derived covariance structure, instead of QTL-derived.
- Representation of covariance structure of the data may be incorrect.
- This may lead to wrong inferences about variance parameters.



Beatriz Castro Dias Cuyabano<sup>1</sup>, Gabriel Rovere<sup>1,2</sup> & Peter Sørensen<sup>1</sup>

- Genomic models have their likelihood misspecified to that of the true model that generates a phenotype.
- Misspecification: SNP-derived covariance structure, instead of QTL-derived.
- Representation of covariance structure of the data may be incorrect.
- This may lead to wrong inferences about variance parameters.
- Sequential statistics:



Beatriz Castro Dias Cuyabano<sup>1</sup>, Gabriel Rovere<sup>1,2</sup> & Peter Sørensen<sup>1</sup>

- Genomic models have their likelihood misspecified to that of the true model that generates a phenotype.
- Misspecification: SNP-derived covariance structure, instead of QTL-derived.
- Representation of covariance structure of the data may be incorrect.
- This may lead to wrong inferences about variance parameters.
- Sequential statistics:
  - ▶ Tool to evaluate potential bias of heritability estimates.



Beatriz Castro Dias Cuyabano<sup>1</sup>, Gabriel Rovere<sup>1,2</sup> & Peter Sørensen<sup>1</sup>

- Genomic models have their likelihood misspecified to that of the true model that generates a phenotype.
- Misspecification: SNP-derived covariance structure, instead of QTL-derived.
- Representation of covariance structure of the data may be incorrect.
- This may lead to wrong inferences about variance parameters.
- Sequential statistics:
  - ▶ Tool to evaluate potential bias of heritability estimates.
  - ▶ Combine information from GBLUP to observed phenotypes.



Beatriz Castro Dias Cuyabano<sup>1</sup>, Gabriel Rovere<sup>1,2</sup> & Peter Sørensen<sup>1</sup>

- Genomic models have their likelihood misspecified to that of the true model that generates a phenotype.
- Misspecification: SNP-derived covariance structure, instead of QTL-derived.
- Representation of covariance structure of the data may be incorrect.
- This may lead to wrong inferences about variance parameters.
- Sequential statistics:
  - ▶ Tool to evaluate potential bias of heritability estimates.
  - ▶ Combine information from GBLUP to observed phenotypes.
- Simulations of several different scenarios (genetic architecture of trait).



Beatriz Castro Dias Cuyabano<sup>1</sup>, Gabriel Rovere<sup>1,2</sup> & Peter Sørensen<sup>1</sup>

